



# QUINCLORAC 75DF SELECT™

HERBICIDE

## *Post-Emergent Weed Control for Turfgrass*



### ACTIVE INGREDIENT

Quinclorac

### % ACTIVE INGREDIENT

18.92%

### FORMULATION

Dry Flowable

### CHEMICAL FAMILY

Quinoline Carboxylic Acid

### MODE OF ACTION

Synthetic Auxin

### RESTRICTED USE

None

### SIGNAL WORD

Caution

### PACKAGING INFORMATION

6 x 1 lb.

### LABELED USES

Residential Turfgrass

Non-Residential Turfgrass

**Quinclorac 75DF Select™** is a selective, post-emergent herbicide that can be safely applied to many types of turf to control a variety of broadleaf and grassy weeds. Quinclorac 75DF Select can be applied to both residential and non-residential turf. Quinclorac 75DF Select can be applied from 7 days before seeding and up to 28 days after emergence on selected weeds. The use of a methylated seed oil, such as Duo Stick Select is recommended to greatly enhance the effectiveness of the product.

- Up to 90 day control
- Works on both broadleaf and grassy weeds
- Selective control means it doesn't harm turf
- Controls Crabgrass
- Use conjunction with seeding/overseeding
- Add Duo Stick Select™ to enhance efficacy

**IMPORTANT:**  
ALWAYS READ AND FOLLOW  
LABEL INSTRUCTIONS.



**PRIME SOURCE**

# QUINCLORAC 75DF SELECT™

HERBICIDE

## General Information:

Quinclorac 75DF Select™ is an auxin agonist and is classified as a quinoline carboxylic acid. It is absorbed by foliage and roots and translocated throughout the plant. The control symptoms exhibited by broadleaf weeds include leaf and stem curl or twisting, and chlorosis. Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death. Apply Quinclorac 75DF Select to actively growing weeds as post-emergence broadcast or spot sprays using the turf species, rate and growth stages indicated on the label.

The use of Quinclorac 75DF Select before or after seeding or overseeding a turf area will not significantly interfere with the turf grass seed germination and growth of those grass types identified as tolerant or moderately tolerant on the label.

## BROADLEAF WEEDS CONTROLLED INCLUDE

Bindweed, field  
Clover, hop  
Clover, red  
Clover, white  
Daisy, English<sup>2,5</sup>  
Dandelion, common<sup>2</sup>  
Dollarweed  
Geranium, Carolina  
Medic, black  
Morningglory spp.  
Speedwell, common  
Speedwell, slender  
Speedwell, thymeleaf  
Violet, wild

## GRASSY WEEDS CON- TROLLED INCLUDE

Barnyardgrass  
Crabgrass, large<sup>1,4</sup>  
Crabgrass, smooth<sup>1,4</sup>  
Foxtail, giant<sup>1</sup>  
Foxtail, green<sup>1</sup>  
Foxtail, yellow<sup>1</sup>  
Kikuyugrass<sup>2,3</sup>  
Signalgrass, broadleaf  
Torpedograss<sup>3</sup>

1. Under certain conditions, application of Quinclorac 75DF Select made to annual grasses at 2 to 4 tiller may not provide complete control. A sequential application will be required for grass control in these situations. Optimum control is achieved when applications of Quinclorac 75DF Select + methylated seed oil are applied either before second tiller or as weed grasses mature.
2. Tank mix partner or sequential application required.
3. Make 2 sequential applications of 1.0 fl. oz. (0.51 lb. AI/A) of Quinclorac 75DF Select per 1,000 sq. ft. and an additional sequential application up to 0.90 fl. oz. (0.46 lb. AI/A) of Quinclorac 75DF Select per 1,000 sq. ft. at 14- to 21-day intervals.
4. Biotypes of large and smooth crabgrass in California have shown varied response to Quinclorac 75DF Select. If control failure occurs following a full or split application, DO NOT reapply Quinclorac 75DF Select. Change to a herbicide with a different mode of action.
5. NOT FOR USE to control this weed in California.



Prime Source, LLC  
4609 E. Boonville-New Harmony Road  
Evansville, IN 47725  
[www.Prime-SourceLLC.com](http://www.Prime-SourceLLC.com)

Download a  
specimen label at  
[Prime-SourceLLC.com](http://Prime-SourceLLC.com)