

TRIAD SELECT™

HERBICIDE

Selective Broad Spectrum Weed Control



ACTIVE INGREDIENTS

2,4-D
MCPA
Dicamba

% ACTIVE INGREDIENT

2,4-D – 30.89%
MCPA – 8.23%
Dicamba – 2.77%

FORMULATION

Liquid Concentrate

CHEMICAL FAMILY

Phenoxyaliphatic Acid,
Benzoic Acid

BRAND ALTERNATIVE

Unique Formulation

RESTRICTED USE

None

SIGNAL WORD

Warning

PACKAGING INFORMATION

4x1 gal.
2x2.5 gal.
30 gal.
55 gal.
250 gal.

Triad Select™ utilizes a combination of three selective herbicides to control a wide variety of hard-to-kill broadleaf weeds such as dandelion, clover and plantain. Triad Select has been demonstrated through university trials to be an effective solution that can be applied to both cool and warm season grasses, and is labeled for use on turf, sod farms, rights-of-ways and other non-crop industrial areas.

- Combined power of three selective herbicides
- For use on both cool and warm season grasses
- Can be tank mixed with liquid fertilizers or liquid iron materials**
- Flexible application rates
- May be applied to woody brush in Non-Cropland areas
- University tested

** Refer to label for fertilizer tank mix compatibility

IMPORTANT:
ALWAYS READ AND FOLLOW
LABEL INSTRUCTIONS.



PRIME SOURCE


TRIAD SELECT™

HERBICIDE

Proven Results:


Virginia Tech - Cool Weather Trial

Applied to: Tall Fescue, Kentucky Bluegrass, Perennial Ryegrass
Application rate = 3pts/acre
16 Days After Treatment



Clemson University - Warm Weather Trial

Applied to: Bermuda, Tall Fescue
Application rate = 4pts/acre
35 Days After Treatment



Application Information:

Bahiagrass, bluegrass, common Bermudagrass, fescue, ryegrass, zoysia grass

Use Rate	Spray Volume	Use Rate	Spray Volume
(Fluid Ounces)	(Gallons)	(Pints)	(Gallons)
Per 1,000 SF		Per Acre	
1.1 to 1.5	.5 to 5	3 to 4	5 to 220

Non-Turf Application Rate

Roadsides (including aprons and guardrails), rights-of-way and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1.5 to 3.5 quarts of product per acre. This may require 50 to 300 gallons of water per acre (43,560 sq ft). For woody plants, apply to both stems and foliage of completely matured plants until they start to go dormant.

Woody Brush: Add 1 to 1.5 gallons of this product per acre to wet all parts of the brush foliage, stem and bark. This may require up to 200 to 600 gallons of water per acre (43,560 sq ft).

See label for complete application rates and recommendations.

Dormant St. Augustine

(Common, Raleigh, and Seville varieties grown in Texas, Louisiana, and Mississippi only)

Use Rate	Spray Volume	Use Rate	Spray Volume
(Fluid Ounces)	(Gallons)	(Pints)	(Gallons)
Per 1,000 SF		Per Acre	
0.75 to 0.90 Maximum	1 to 5	2 to 2.5 Maximum	40 to 200

Bentgrass

(Putting and bowling greens)

Use Rate	Spray Volume	Use Rate	Spray Volume
(Fluid Ounces)	(Gallons)	(Pints)	(Gallons)
Per 1,000 SF		Per Acre	
0.67 Maximum	3.3	1.8 Maximum	145

WEEDS CONTROLLED

Bedstraw
Black Medic
Buckhorn
Chicory
Chickweed
Clover
Dandelion
Dock
Ground Ivy
Heal-All
Henbit
Knotweed
Lambsquarters
Lespedeza
Mallow
Morningglory
Peppergrass
Pigweed
Plantain
Poison Ivy
Poison Oak
Purslane
Ragweed
Sheep Sorrel
Shepherdspurse
Speedwell
Spurge
Wild Carrot
Wild Garlic
Wild Lettuce
Wild Onion
Yarrow & other broadleaf weeds

APPLICATION AREAS

Golf Courses
Cemeteries
Parks & Sports Fields
Turfgrass, Lawns & Grass
Turf Grown for Seed & Sod
Fence & Hedge Rows
Roadsides & Ditches
Rights-of-Way
Utility Power Lines
Railroads, Airports & Industrial Sites

Download a
specimen label at
Prime-SourceLLC.com



Prime Source, LLC
4609 E. Boonville-New Harmony Road
Evansville, IN 47725
www.Prime-SourceLLC.com