



QUINCLORAC 1.5L SELECT™

HERBICIDE

Post-Emergent Weed Control for Turfgrass



ACTIVE INGREDIENT

Quinclorac

% ACTIVE INGREDIENT

18.92%

FORMULATION

Flowable Concentrate

CHEMICAL FAMILY

Quinoline Carboxylic Acid

MODE OF ACTION

Synthetic Auxin

RESTRICTED USE

None

SIGNAL WORD

Caution

PACKAGING INFORMATION

4 x 0.5 gal.

LABELED USES

Residential Turfgrass

Non-Residential Turfgrass

Quinclorac 1.5L Select™ is a selective, post-emergent herbicide that can be safely applied to many types of turf to control a variety of broadleaf and grassy weeds. Quinclorac 1.5L Select can be applied to both residential and non-residential turf. Quinclorac 1.5L Select can be applied from 7 days before seeding and up to 28 days after emergence on selected weeds. The use of a methylated seed oil, such as Duo Stick Select is recommended to greatly enhance the effectiveness of the product.

- Up to 90 day control
- Works on both broadleaf and grassy weeds
- Selective control means it doesn't harm turf
- Controls Crabgrass
- Use conjunction with seeding/overseeding
- Add Duo Stick Select™ to enhance efficacy

IMPORTANT:
ALWAYS READ AND FOLLOW
LABEL INSTRUCTIONS.


PRIME SOURCE

QUINCLORAC 1.5L SELECT™

HERBICIDE

General Information:

Quinclorac 1.5L Select is an auxin agonist and is classified as a quinoline carboxylic acid. It is absorbed by foliage and roots and translocated throughout the plant. The control symptoms exhibited by broadleaf weeds include leaf and stem curl or twisting, and chlorosis. Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death. Apply Quinclorac 1.5L Select to actively growing weeds as post-emergence broadcast or spot sprays using the turf species, rate and growth stages indicated on the label.

The use of Quinclorac 1.5L Select before or after seeding or overseeding a turf area will not significantly interfere with the turf grass seed germination and growth of those grass types identified as tolerant or moderately tolerant on the label.

BROADLEAF WEEDS CONTROLLED INCLUDE

Bindweed, field
Clover, hop
Clover, red
Clover, white
Daisy, English^{2,5}
Dandelion, common²
Dollarweed
Geranium, Carolina
Medic, black
Morningglory spp.
Speedwell, common
Speedwell, slender
Speedwell, thymeleaf
Violet, wild

GRASSY WEEDS CON- TROLLED INCLUDE

Barnyardgrass
Crabgrass, large^{1,4}
Crabgrass, smooth^{1,4}
Foxtail, giant¹
Foxtail, green¹
Foxtail, yellow¹
Kikuyugrass^{2,3}
Signalgrass, broadleaf
Torpedograss³

1. Under certain conditions, application of Quinclorac 1.5L Select made to annual grasses at 2 to 4 tiller may not provide complete control. A sequential application will be required for grass control in these situations. Optimum control is achieved when applications of Quinclorac 1.5L Select + methylated seed oil are applied either before second tiller or as weed grasses mature.
2. Tank mix partner or sequential application required.
3. Make 2 sequential applications of 1.0 fl. oz. (0.51 lb. AI/A) of Quinclorac 1.5L Select per 1,000 sq. ft. and an additional sequential application up to 0.90 fl. oz. (0.46 lb. AI/A) of Quinclorac 1.5L Select per 1,000 sq. ft. at 14- to 21-day intervals.
4. Biotypes of large and smooth crabgrass in California have shown varied response to Quinclorac 1.5L Select. If control failure occurs following a full or split application, DO NOT reapply Quinclorac 1.5L Select. Change to a herbicide with a different mode of action.
5. NOT FOR USE to control this weed in California.



Prime Source, LLC
4609 E. Boonville-New Harmony Road
Evansville, IN 47725
www.Prime-SourceLLC.com

Download a
specimen label at
Prime-SourceLLC.com