



**PRIME SOURCE**

# CROSSCUT SELECT™

*Post-emergent brush, woody plant and broadleaf weed control*

## ACTIVE INGREDIENTS

Triclopyr  
2,4-D ester

## % ACTIVE INGREDIENTS

2,4-dichlorophenoxyacetic acid, butoxyethyl ester. . . . . 34.4%  
Triclopyr BEE: 3,5,6-trichloro-2-pyridinyloxyacetic acid, butoxyethyl ester . . 16.5%

## FORMULATION

Liquid

## CHEMICAL FAMILY

Chlorinated  
phenoxy-systemic herbicide

## BRAND ALTERNATIVE

Crossbow®

## SIGNAL WORD

Caution

## PACKAGING

12 x 1 qt, 4 x 1 gal  
2 x 2.5 gal

## LABELED USES

Commercial and recreational turf  
Commercial sod  
Rangeland  
Permanent grass pastures  
Conservation reserve program (CRP) acres  
Fence rows  
Non-irrigation ditchbanks  
Roadsides and other non-crop areas  
Industrial sites

## IMPORTANT:

ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS.



**CrossCut Select™** is a post-emergent herbicide that provides control of most woody plants and brush, as well as annual and perennial broadleaf weeds on rangeland, permanent grass pastures, recreational turf, commercial turf, sod farms, conservation reserve program (CRP) acres, fence rows, non-irrigation ditch banks, roadsides, other non-crop areas and industrial sites. Formulated as a low volatile ester, the combination of triclopyr and 2,4-D provides outstanding weed control in both cool and warm weather.

- Provides excellent control of brush and woody plants
- Controls broadleaf weeds with no harm to turf
- Low volatility formulation
- Can be applied by all types of sprayers
- Rainfast within 2 hours post application
- Flexible application timing – can be applied in-season or during dormant periods
- Apply to actively growing foliage for best results

## FOR USE ON TURF

### Broadcast treatment of commercial and recreational turf and commercial sod farms

To be applied only under the direct supervision of Commercial applicators responsible for turf weed control programs.

Apply 2 to 4 pints (.5 to 1 lb. 2,4-D ai) of **CrossCut Select** in enough water to make 20 to 200 gallons total spray per acre to control broadleaf weeds growing in tall fescue, bluegrass, or perennial ryegrass turf. Do not use on other grass species, such as bentgrass or St. Augustine grass, unless injury can be tolerated.

### RESTRICTIONS: For broadcast application on commercial, and recreational turf

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

### Post-emergence:

- Limited to 2 applications per year.
- Use a maximum of 4 pints **CrossCut Select** (1.5 lbs. 2,4-D ai) per acre per application.
- The maximum seasonal rate is 8 pints **CrossCut Select** per acre (2.0 lbs. 2,4-D ai/A), excluding spot treatments.

# CROSSCUT SELECT™

Post-emergent brush, woody plant and broadleaf weed control

## FOR USE ON RANGELAND AND PASTURE

### Broadcast treatment (ground equipment and helicopter):

Apply up to 4 quarts of **CrossCut Select** per acre in sufficient water to deliver 10 to 30 gallons of total spray per acre. Actively growing weeds are most susceptible. Optimal application time for biennial and winter annual weeds is the rosette stage. Hard to control weeds such as field bindweed, chicory, dogfennel, goldenrod, horsenettle, kudzu, milkweed, perennial sowthistle, leafy spurge, and Canada thistle may require retreatment for complete control.

### Spot Treatment:

To control broadleaf weeds in small areas with a hand sprayer, mix 4 to 6 fl. oz. of **CrossCut Select** in 3 gallons of water. Spray to thoroughly wet all foliage.

### RESTRICTIONS For rangeland and pasture

- Do not apply more than 4 quarts per acre per application.
- Do not make more than one application per year.
- Pre-harvest interval is 14 days
- Except for lactating dairy animals there are no grazing restrictions
- Do not apply more than 1 lb. 2,4-D ai per application (2 qts. product) for susceptible annual and biennial broadleaf weeds.

### Weeds controlled include:

Black medic	Chickweeds	Field Bindweed	Poison Ivy
Bull thistle	(common and mouse ear)	Ground Ivy	Poison Oak
Burdock	Cinquefoil	Henbit	Speedwell
Buttercup	Clovers	Lambsquarters	Smartweed
Blackberry	Creeping beggar weed	Oxalis (stricta and corniculata)	Vetch
Canada thistle	Dandelion	Plantain	Wild violet
	Docks		

### Woody plant control (easy to control species):

6 quarts/acre broadcast application or 1 to 1.5% mixtures for high-volume foliar applications.

Alder	Dogwood	Sumac	Speedwell
Ash	Elderberry	Sycamore	Smartweed
Beech	Hawthorn	Tamarack	Vetch
Blackberry Cherry (except black)	Poison ivy	Wax myrtle (top growth)	Wild violet
Cottonwood	Poison oak	White oak	And more
	Scotch broom	Wild grape	

### Woody plant control (harder to control species):

Apply high-volume applications, 1.5% mixture, conventional basal or dormant stem applications. A broadcast rate of 8 quarts/A will increase the degree of control of these species.

Buckbush (Symphoricarpos spp.) (suppression)	Hazel	Salmonberry (suppression)
Common persimmon (suppression)	Honeylocust (suppression)	Sweetgum
Elm (except winged elm)	Pine (suppression)	Trumpet creeper (suppression)
	Russian olive	Virginia creeper

Refer to label for full list of broadleaf and grassy weeds controlled.