



PRIME SOURCE

# TRIAD **TZ** SELECT™

Broad spectrum weed control

## ACTIVE INGREDIENTS

2,4-D, 2-ethylhexyl ester  
Triclopyr BEE  
Dicamba acid  
Sulfentrazone

## % ACTIVE INGREDIENTS

2,4-D, 2-ethylhexyl ester  
..... 31.82%  
Triclopyr BEE ..... 8.40%  
Dicamba acid ..... 2.43%  
Sulfentrazone. .... 0.73%

## FORMULATION

Emulsifiable Concentrate

## CHEMICAL FAMILY

PPO inhibitor (Group 14),  
Growth Regulator (Group 4)

## BRAND ALTERNATIVE

T-Zone™

## SIGNAL WORD

Caution

## PACKAGING

12 x 1 qt, 4 x 1 gal,  
2 x 2.5 gal, 5 x 30 gal

## LABELED USES

Residential lawns  
Athletic fields  
Golf courses  
Parks  
Industrial and  
institutional buildings  
Commercial sod production  
Fencerows  
Farmyards  
Utility rights-of-way

## IMPORTANT:

ALWAYS READ  
AND FOLLOW LABEL  
INSTRUCTIONS.



**Triad TZ Select™** contains four active ingredients, with two modes of action to provide dependable post-emergent control of broadleaf weeds. It controls the listed annual and perennial broadleaf weeds in tall and fine fescues, Kentucky bluegrass, and perennial ryegrass. **Triad TZ Select™** may also be applied to fully dormant bermudagrass, fully dormant zoysiagrass and fully dormant bahiagrass. **Triad TZ Select™** is quickly taken up by the leaves and stems of plants and weed growth ceases within hours. The weed control effects may be seen within a few hours to a few days after application. Weed death occurs in 1 to 4 weeks after application, depending on the application rate, sensitivity of the weed, and environmental conditions.

- Excellent post-emergent activity with proven rapid and effective weed control performance for some of the toughest broadleaf weeds in turfgrass such as dandelion, spurge, white clover, plantains, ground ivy, Virginia buttonweed and wild violet. Labelled for the suppression of yellow nutsedge.
- This product exhibits improved cool-weather performance.
- The speed of action (rate of weed phytotoxicity, yellowing) and the early weed symptoms are features of sulfentrazone. Often, the weed injury symptoms can be noticed within hours of the application and plant death can occur within 10 to 14 days.
- This product is generally rainfast in as little as three hours.

Prime Source, LLC • 10025 US 264 Alternate, Middlesex, NC 27557 • [www.prime-sourcellc.com](http://www.prime-sourcellc.com)

©2019 Prime Source, LLC. Select is a trademark of Excelsior Equity, T-Zone is a trademark of PBI-Gordon Corporation. Always read and follow label directions for use.

# TRIAD T<sub>Z</sub> SELECT™

Broad spectrum weed control

## USE RATES AND SPRAY VOLUMES:

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

**Table 1. Use rates for ornamental turfgrass, sod farms and non-cropland.**

Species	Rate	Spray Volume
<b>Cool season turf</b>		
Kentucky bluegrass, annual bluegrass, annual ryegrass perennial ryegrass, tall fescue, red or fine leaf fescues	3.25 to 4 pints/A (1.2 to 1.5 fl. oz./1,000 sq.ft.)	10 to 220 gal/A (0.25 to 5.0 gal/1,000 sq.ft.)
<b>Warm season turf (Dormant turf)</b>		
Hybrid Bermudagrass, common Bermudagrass, Zoysiagrass and Bahiagrass	2 to 2.25 pints/A (0.75 to 0.83 fl. oz./1,000 sq.ft.)	10 to 220 gal/A (0.25 to 5.0 gal/1,000 sq.ft.)
<p><b>Dormant turf:</b> This product may be applied to fully dormant bermudagrass, fully dormant zoysiagrass and fully dormant bahiagrass.</p> <p><b>Note:</b> Do not apply to above listed warm-season turfgrass unless turf injury can be tolerated. It is impossible to test all environmental conditions for the listed warm-season turfgrass. We suggest testing this product on a small area and observe the treated area for 30 days to determine the acceptability of turf discoloration.</p> <p>Do not apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy.</p>		

**TABLE 2. Limitations on broadcast applications to ornamental turfgrass, sod farms and non-cropland.**

Use Site	Maximum rate per application	Maximum number of applications per year*	Minimum interval between applications	Maximum seasonal rate
Ornamental turfgrass	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	2	N/A	8.0 pints/A (1.75 lb 2,4-D ae/A) (0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A)
Sod farms	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb. triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	2	21 days	8.0 pints/A (1.75 lb 2,4-D ae/A) (0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A)
Non-cropland	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	1 (woody plants)	N/A	8.0 pints/A (1.75 lb 2,4-D ae/A) (0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A)
		2 (annual & perennial weeds)	30 days	

\* New York: Only one application per year of this product is allowed.

Refer to label for full list of broadleaf and grassy weeds controlled.