



PRIME SOURCE

TRIAD SELECT™

Selective broad spectrum weed control

ACTIVE INGREDIENTS

2,4-D
MCPA
Dicamba

% ACTIVE INGREDIENTS

2,4-D 30.89%
MCPA 8.23%
Dicamba 2.77%

FORMULATION

Liquid concentrate

CHEMICAL FAMILY

Phenoxyaliphatic acid,
Benzoic acid

BRAND ALTERNATIVE

Unique formulation

RESTRICTED USE

None

SIGNAL WORD

Warning

PACKAGING

4 x 1 gal, 2 x 2.5 gal, 30 gal,
55 gal, 265 gal

LABELED USES

Golf courses
Cemeteries
Parks & sports fields
Turfgrass, lawns & grass
Turf grown for seed & sod
Fence & hedgerows
Roadsides & ditches
Rights-of-way
Utility power lines
Railroads, airports &
industrial sites

IMPORTANT:

**ALWAYS READ
AND FOLLOW LABEL
INSTRUCTIONS.**



Triad Select™ utilizes a combination of three selective herbicides to control a wide variety of hard-to-kill broadleaf weeds such as dandelion, clover and plantain. **Triad Select** has been demonstrated through university trials to be an effective solution that can be applied to both cool and warm season grasses, and is labeled for use on turf, sod farms, rights-of-ways and other non-crop industrial areas.

- Combined power of three selective herbicides
- Low odor formulation with flexible application rates
- For use on both cool and warm season grasses
- May be applied to woody brush in Non-Cropland areas
- Can be tank mixed with liquid fertilizers or liquid iron materials*
- University tested

*Refer to label for fertilizer tank mix compatibility.

Prime Source, LLC • 10025 US 264 Alternate, Middlesex, NC 27557 • www.prime-sourcellc.com

©2019 Prime Source. Select™ is a trademark of Prime Source LLC. Trimec® is a registered trademark of PBI/Gordon Corporation.

TRIAD SELECT™


Selective broad spectrum weed control

Proven Results

Virginia Tech - Cool Weather Trial

% Percent control @ 16 DAT

Applied to Tall fescue, Kentucky bluegrass and Perennial ryegrass • Untreated = 0% Control



Clemson University - Warm Weather Trial

% Percent control @ 35 DAT

Applied to: Bermuda, Tall fescue
Untreated = 0% Control



Application information

Bahiagrass, bluegrass, common bermudagrass, fescue, ryegrass, zoysia grass

Use Rate Fluid Ounces	Spray Volume Gallons	Use Rate Fluid Ounces	Spray Volume Gallons
Per 1,000 sf		Per Acre	
1.1 to 1.5	.5 to 5	3 to 4	5 to 220

Dormant St. Augustine

(Common, Raleigh, and Seville varieties grown in Texas, Louisiana, and Mississippi only)

Use Rate Fluid Ounces	Spray Volume Gallons	Use Rate Fluid Ounces	Spray Volume Gallons
Per 1,000 sf		Per Acre	
0.75 to 0.90 maximum	1 to 5	2 to 2.5 maximum	40 to 200

Bentgrass

(Putting and bowling greens)

Use Rate Fluid Ounces	Spray Volume Gallons	Use Rate Fluid Ounces	Spray Volume Gallons
Per 1,000 sf		Per Acre	
0.67 maximum	3.3	1.8 maximum	145

Refer to label for full list of broadleaf and grassy weeds controlled.

Non-turf application rate

Roadsides (including aprons and guardrails), rights-of-way and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1.5 to 3.5 quarts of product per acre. This may require 50 to 300 gallons of water per acre (43,560 sq ft). For woody plants, apply to both stems and foliage of completely matured plants until they start to go dormant.

Woody Brush: Add 1 to 1.5 gallons of this product per acre to wet all parts of the brush foliage, stem and bark. This may require up to 200 to 600 gallons of water per acre (43,560 sq ft).

Weeds controlled include:

Bedstraw	Knotweed	Ragweed
Black Medic	Lambsquarters	Sheep Sorrel
Buckhorn	Lespedeza	Shepherdspurse
Chicory	Mallow	Speedwell
Chickweed	Morningglory	Spurge
Clover	Peppergrass	Wild Carrot
Dandelion	Pigweed	Wild Garlic
Dock	Plantain	Wild Lettuce
Ground Ivy	Poison Ivy	Wild Onion
Heal-All	Poison Oak	Yarrow & other broadleaf weeds
Henbit	Purslane	