

AZOXY 2SC SELECT[™]

Long term protection in a convenient liquid formulation

ACTIVE INGREDIENT

Azoxystrobin – Group 11

% ACTIVE INGREDIENT

Azoxystrobin 22.9%

FORMULATION

Suspended concentrate

CHEMICAL FAMILY

Strobilurin/QoI inhibitors

BRAND ALTERNATIVE

Heritage[™] TL

MODE OF ACTION

Acropetal penetrant

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

12 x 1 pt, 4 x 1 gal

LABELLED USES

- Golf turf
- Residential turf
- Commercial turf
- Parks/Rec
- Athletic fields



Azoxyl 2SC Select[™] is the equivalent liquid formulation to the popular WDG azoxystrobin products. The convenient suspended concentrate formulation allows for convenient tank mixing and application. Total protection is achieved through a combination of foliar and root absorption. Highly systemic and with a higher than average degree of root uptake, **Azoxyl 2SC Select[™]** provides continuous protection to the whole plant for up to 28 days.

PRODUCT BENEFITS

- Convenient liquid formulation for easy tank mixing
- Defends against both soil borne and foliar diseases
- Complete root-to-leaf distribution
- Constant delivery for even distribution to new growth
- Greater degree of root uptake when compared to other strobilurins
- Prevents spore germination, thereby preventing many fungal infections
- Does not impact honey bees and other beneficials
- Low use rate

IMPORTANT: ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS.

AZOXY 2SC SELECT™

Long term protection in a convenient liquid formulation

APPLICATION INFORMATION

Azoxy 2SC Select™ may be applied as a foliar spray, in alternating spray programs, or in tank mixes with other registered pesticides.

Azoxy 2SC Select™ is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides, and biological control products. If tank mixes are desired, observe all directions, precautions, and limitations on labeling of all products used.

Azoxy 2SC Select™ is recommended for control of anthracnose, brown patch, cool weather brown patch (yellow patch), *Fusarium* patch, gray leaf spot, gray snow mold (*Typhula* blight), leafspot, melting out, necrotic ring spot, pink patch, pink snow mold, *Pythium* blight, *Pythium* root rot, red thread, *Rhizoctonia* large patch, southern blight, spring dead spot, summer patch, take-all patch, and *Zoysia* patch on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Azoxy 2SC Select™ is most effective when applied prior to disease development. Mix **Azoxy 2SC Select™** with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1000 square feet (87-174 gallons per acre). Repeat applications at specified intervals for as long as required. For spot treatments, use 0.4 oz. **Azoxy 2SC Select™** per 1 to 2 gallons of water. Do not apply more than 9.6 quarts product/acre/year (7.1 fl. oz. product/1000 square feet/year). Make applications by ground only.

Labeled Diseases

Anthracnose	Leaf spot	<i>Pythium</i> root rot	Take-All patch
Brown patch	Melting out	Red thread	<i>Typhula</i> blight
Cool weather brown patch	Necrotic ring spot	<i>Rhizoctonia</i> large patch	Yellow patch
<i>Fusarium</i> patch	Pink patch	Spring dead spot	<i>Zoysia</i> patch
Gray leaf spot	Pink snow mold	Southern blight	
Gray snow mold	<i>Pythium</i> blight	Summer patch	