

FLUAZINAM 40SC SELECT[™]

Broad spectrum turf fungicide

ACTIVE INGREDIENT

Fluazinam – Group 29

% ACTIVE INGREDIENT

Fluazinam40%

FORMULATION

Suspension concentrate

CHEMICAL FAMILY

2,6-dinitroanilines

BRAND ALTERNATIVE

Secure®

MODE OF ACTION

Oxidative phosphorylation
uncoupler

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

4 x 0.5 gal, 2 x 2.5 gal

LABELED USES

Golf course tees, greens,
fairways and roughs



Fluazinam 40SC Select[™] is a broad spectrum, multi-site fungicide to control diseases in turf grasses on golf courses only. The active ingredient, fluazinam, is the only registered fungicide for turf in FRAC 29 group. It has no known resistance and as a multi-site contact, it has a very low risk of developing resistance. When partnered with chlorothalonil-based fungicides, **Fluazinam 40SC Select**[™] provides season-long protection against the toughest turf diseases without interruptions.

PRODUCT BENEFITS

- Unique multi-site contact active ingredient, reduces resistance management
- Excellent control of dollar spot and 10 additional turf diseases
- Perfect tank mix partner with systemics for inside out protection
- Controls dollar spot that is resistant to other chemistries

**IMPORTANT: ALWAYS
READ AND FOLLOW
LABEL INSTRUCTIONS.**

FLUAZINAM 40SC SELECT™

Broad spectrum turf fungicide

APPLICATION INFORMATION

Apply **Fluazinam 40SC Select™** in sufficient water to obtain adequate uniform coverage of the turf. Apply in a minimum spray volume of 40 gal of water per acre on fairways and 90 gal of water per acre on tees and greens.

- Apply at 0.5 fl. oz. per 1,000 sq. ft. (6 acres per gallon) every 14 days.
- Apply after mowing with sufficient water to uniformly cover the turf and allow area to dry completely before irrigation.
- Refer to the product label for additional information about this product and tank mixing.
- Not for use on home lawns or any residential turf.

Turfgrass disease control

- Use **Fluazinam 40SC Select™** in a preventative disease control program.
- Apply in sufficient water to ensure thorough coverage.
- Apply after mowing or allow sprayed area to completely dry before mowing.
- For control of foliar diseases, allow sprayed area to completely dry before irrigation.
- For optimum turf quality and disease control, use **Fluazinam 40SC Select™** in conjunction with turf management practices that promote good plant health and optimum disease control.
- Evaluate spray additives prior to use. Label directions are based on data obtained with no additives.
- Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.



Restrictions

- **DO NOT** exceed 258 fl oz of **Fluazinam 40SC Select™** (8.4 pounds active ingredient) per acre, per year.
- **DO NOT** apply more than twelve (12) applications at 21.5 fl oz per acre (0.5 fl oz/1000 sq ft), per year.

Labeled diseases

Algal scum	Gray snow mold	Pink patch
Anthracnose	Large patch	Pink snow mold
Brown blight	Leaf spot	Red thread
Brown patch	Melting out	Stem rust
Dollar spot		

Read label for full application and rate information.

Prime Source, a Division of Albaugh, LLC • 1525 NE 36th Street, Ankeny, IA 50021 • www.prime-sourcellc.com

©2021 Albaugh, LLC. Select is a trademark of Albaugh, LLC. Secure is a registered trademark of Syngenta Group Company.