

TM 4.5F SELECT[™]

Broad spectrum systemic fungicide

ACTIVE INGREDIENT

Thiophanate methyl –
Group 1

% ACTIVE INGREDIENT

Thiophanate methyl. .46.2%

FORMULATION

Flowable

CHEMICAL FAMILY

Thiophanates

BRAND ALTERNATIVE

Cleary's 3336[®] F

Topsin[®] 4.5FL

MODE OF ACTION

Systemic
(mitosis and cell division)

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

2 x 2.5 gal

LABELED USES

Greenhouse

Nursery

Golf courses

Athletic fields

Cemeteries

Parks

Commercial lawns

Residential lawns

Landscape ornamentals

Conifers*

*not for use on conifers
in California



TM 4.5F Select[™] is a broad spectrum, systemic fungicide for the control and prevention of diseases on ornamentals and cool and warm season turfgrasses. **TM 4.5F Select[™]** is the perfect disease management tool to keep your golf course or athletic field in peak condition. For optimal effect in areas that utilize irrigation, add a spreader/sticker adjuvant to the tank mix.

PRODUCT BENEFITS

- No known phytotoxicity to turfgrass
- Both preventive and curative properties
- Use on both cool and warm season turfgrasses
- Can be used through certain irrigation systems

**IMPORTANT: ALWAYS
READ AND FOLLOW
LABEL INSTRUCTIONS.**

Prime Source, a Division of Albaugh, LLC • 1525 NE 36th Street, Ankeny, IA 50021 • www.prime-sourcellc.com

©2021 Albaugh, LLC. Select is a trademark of Albaugh, LLC. Cleary's 3336 is a registered trademark of Cleary Chemicals LLC. Topsin is a registered trademark of Nippon Soda Company, Ltd.

TM 4.5F SELECT™

Broad spectrum, systemic fungicide

APPLICATION INFORMATION

Horticultural applications

TM 4.5F Select™ provides broadspectrum control of many foliar, stem, and below-ground diseases on a wide range of horticultural plants grown or maintained under a variety of conditions. Apply TM 4.5F Select™ 14 to 21 days prior to when a particular disease usually appears and at the very latest, upon first sign of disease. Use 7- to 14-day spray intervals with 14 days being for preventive treatments and the 7-day interval for times when conditions are conducive to disease development. Add an acceptable wetting agent to the spray tank to increase product efficacy for hard-to-wet foliage. Use a spreader/sticker when excessive and repeated foliar wetting occurs.

Turf applications

Use TM 4.5F Select™ against certain foliar and soil diseases for use on all turf types such as golf course greens, tees and fairways, athletic fields, cemeteries, parks, and commercial and home lawns. Use TM 4.5F Select™ both preventively and curatively; it is not phytotoxic on turfgrass. Do not use TM 4.5F Select™ on turf being grown for sale or other commercial uses such as sod. For use only by certified applicators and those under their direct supervision. Do not apply with fixed wing or rotary aircraft. Use TM 4.5F Select™ on all cool and warm season grasses (such as but not limited to bentgrasses, bermudagrasses, bluegrasses, fescues, ryegrasses, St. Augustinegrasses and Zoysiagrasses) or their mixtures.

Mixing instructions

Add the required amount of TM 4.5F Select™ to a partially filled tank agitated by mechanical or hydraulic means. Add the remaining required amount of water. Maintain continuous agitation to keep the material in suspension. For best results, use spray mixture the same day it is prepared.

Labeled diseases

Anthracnose	Zoysia patch	Summer patch	Stripe smut
Dollar spot	Red thread	<i>Fusarium</i> blight	
Copper spot	Pink snow mold	Necrotic ring spot	
Brown patch	Gray leaf spot	Spring dead spot	

