

TEBUCONAZOLE 3.6 SELECT[™]

Curative and preventative fungicide

ACTIVE INGREDIENTS

Tebuconazole – Group 3

% ACTIVE INGREDIENT

Tebuconazole 38.7%

FORMULATION

Suspension concentrate

CHEMICAL FAMILY

Triazole

BRAND ALTERNATIVE

Torque[™]

MODE OF ACTION

Demethylation inhibitor

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

4 x 1 gal

LABELLED USES

Golf course turf only

Nursery ornamentals

Commercial

landscape ornamentals

Residential

landscape ornamentals



Tebuconazole 3.6 Select[™] provides long-lasting curative and preventative disease control to protect your ornamentals and golf course turf. Effective against mildews, rusts, blights and many more difficult to control diseases.

PRODUCT BENEFITS

- Preventive and curative properties
- Provides control up to 21 days
- Acropetal (upward) movement
- Rapid uptake (4 hours)
- Highly compatible tank mix partner
- Low use rates

**IMPORTANT: ALWAYS
READ AND FOLLOW
LABEL INSTRUCTIONS.**

TEBUCONAZOLE 3.6 SELECT™

Curative and preventative fungicide

APPLICATION INFORMATION

Turf

For use on all golf turf applications of cool season and warm season grasses (such as bentgrasses, bluegrasses, fescues, ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. **Tebuconazole 3.6 Select™** is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to **Tebuconazole 3.6 Select™** under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85°F.

Restrictions

- For use on golf course turf only.
- **DO NOT** use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- Not for homeowner use.
- Not for use on turf being grown for sale or commercial use as sod.
- **DO NOT** use clippings for animal feed.
- **DO NOT** exceed 3.6 fl. oz. of **Tebuconazole 3.6 Select™** per 1,000 sq. ft. per year.
- **DO NOT** apply more than 6 applications/year except in New York. In New York State, do not apply more than 3 applications/year.

Field, nursery and container ornamentals and commercial and residential landscape applications:

Tebuconazole 3.6 Select™ can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when **Tebuconazole 3.6 Select™** is used in conjunction with sound disease management practices. Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease.

Restrictions

- For use on ornamental plants only; not for use on woodlands or forest management.
- Not for homeowner use.
- Do not apply more than 10 fl. oz. per acre in a single application.
- Do not apply more than 0.31 gallons (40 fl. oz.) of **Tebuconazole 3.6 Select™** (equal to 1.13 lbs. of \ tebuconazole) per acre per year.
- Do not make more than 4 applications per year at highest rate.
- Do not apply to bearing fruit trees or vegetables.

Labeled turf diseases

Anthraxnose	Gray snow mold
Bermuda grass decline	Large patch
Brown patch	<i>Microdochium</i> patch
Brown ring patch	Necrotic ring spot
Copper spot	Pink patch
Corticium red thread	Pink snow mold
Dollar spot	Powdery mildew
Fusarium patch	Red thread
Gray leaf spot	Rhizoctonia blight

Labeled ornamental diseases

Anthraxnose
Black spot
Leaf spot
Petal blight
Powdery mildew
Rusts
<i>Rhizoctonia</i> stem and root rot
Southern blight
Tulip fire

Read label for full application and rate information.